

1. Bounded extension of operators. 1. A separable case.

The isomorphic analogue of the class $\mathfrak{P}_1(X^f)$ is the class $\mathfrak{P}^\sim(X^f)$, which consists of all such spaces $E <_f X$, whose *isomorphic* image in every space from X^f admits a bounded projection.

This class essentially differs from $\mathfrak{P}_1(X^f)$ but has much common with $\mathfrak{J}(X^f)$. Below it will be shown that (in difference to the classical case $\mathfrak{P}_1(c_0^f)$) there are such classes X^f that $\mathfrak{P}^\sim(X^f) = \emptyset$ and such classes Y^f that $\mathfrak{P}^\sim(Y^f) \neq \emptyset$ but $\mathfrak{P}_1(X^f) \not\subseteq \mathfrak{P}^\sim(X^f)$.

It is more natural to study the (formally more wide) class $\mathfrak{P}^\sim(X^F)$, define below.

DEFINITION 1. *Let X be a Banach space, X^F be the corresponding class of crudely finite equivalence. A space X_0 is said to be boundedly universally complemented in X^F provided every space $Z \in X^F$, which contains a subspace isometric to X_0 , admits a bounded projection $P : Z \rightarrow X_0$.*

Surely, this definition is equivalent to the formally stronger property of existence of a bounded projection $P' : Z \rightarrow iX_0$ where $i : X_0 \rightarrow Z$ is an isomorphic embedding.

Certainly, every finite-dimensional Banach space A generates the class A^F , which consists of all finite-dimensional Banach spaces B of dimension $\dim B = \dim A = n$.

Since for every finite-dimensional Banach space A and every space Y that contains a subspace A there exists a projection $P : Y \rightarrow A$ of norm $\|P\| \leq \sqrt{\dim A}$ (cf. [25]), all finite-dimensional Banach spaces are universally complemented in each class X^F , which is generated by an infinite-dimensional space X . So, such a case will be eliminated as trivial.

A class of all infinite-dimensional Banach spaces, which are boundedly universally complemented in the class X^F will be denoted by $\mathfrak{P}^\sim(X^F)$.

Similarly to the classes $\mathcal{A}(X^f)$, $\mathcal{A}_0(X^f)$ and $\mathcal{E}(X^f)$ there may be defined their isomorphic analogues – classes $\mathcal{A}^\sim(X^F)$, $\mathcal{A}_0^\sim(X^F)$ and $\mathcal{E}^\sim(X^F)$.

DEFINITION 2. *The class $\mathcal{A}^\sim(X^F)$ consists of those (infinitely-dimensional) Banach spaces X_0 that have the property:*

- *If $Z \in X^F$ and $X_0 \hookrightarrow Z$ then the annihilator $(X_0)^\perp \hookrightarrow Z^*$ is a complemented subspace of Z^* .*

The class $\mathcal{A}_0^\sim(X^F)$ is given by $\mathcal{A}_0^\sim(X^F) = \mathcal{A}^\sim(X^F) \cap X^F$.

DEFINITION 3. *Let $Z \in \mathcal{B}$; $W \hookrightarrow Z$; $\lambda < \infty$. W is said to be a λ -reflecting subspace of Z if for every finitely dimensional subspace $A \hookrightarrow Z$ there exists an isomorphic embedding $u : A \rightarrow W$ with $\|u\| \|u^{-1}\| \leq \lambda$, such that $u|_{A \cap Z} = \text{Id}_{A \cap Z}$.*

Certainly if W is a λ -reflecting subspace of Z for all $\lambda > 1$ then $W \prec_u Z$.

DEFINITION 4. *The class $\mathcal{E}^\sim(X^F)$ consists of those (infinitely-dimensional) Banach spaces W that have the property:*

- *If $Z \in X^F$ and W is isomorphic to a subspace iW of Z then there exists a such $\lambda < \infty$ that the image iW is a λ -reflecting subspace of Z .*

The following theorem is proved similarly to the isometric analogue. Here $\mathfrak{q}\mathfrak{d}^\sim$ denotes the class of all Banach spaces X that have a complement in the second dual X^{**} (under the canonical embedding $k_X : X \rightarrow X^{**}$).

THEOREM 1. $\mathfrak{P}^\sim(X^F) = \mathcal{A}^\sim(X^F) \cap \mathfrak{q}\mathfrak{d}^\sim$.
 If $Y \in \mathcal{A}^\sim(X^F)$ then $Y^{**} \in \mathfrak{P}^\sim(X^F)$.
 If X is superreflexive then $\mathfrak{P}^\sim(X^F) = \mathcal{A}^\sim(X^F)$.

This theorem is of restricted interest.

Indeed, in difference with the isometric case, there no analogue to the non-emptiness of the class $\mathcal{E}^\sim(X^F) \subseteq \mathcal{A}^\sim(X^F)$. Moreover, in some cases even the widest class $\mathcal{A}_0^\sim(X^F)$ also may be empty.

EXAMPLE 1. If $1 \leq p < 2$ then $\mathcal{A}_0^\sim((l_p)^F) = \mathcal{A}^\sim((l_p)^F) = \emptyset$.
 If $2 < p < \infty$ then $\mathcal{A}_0^\sim((l_p)^F) = \emptyset$; $\mathcal{A}^\sim((l_p)^F) = (l_2)^F$.

PROOF. Notice that every space L_p ($1 \leq p \neq 2 < \infty$) contains an uncomplemented space, isomorphic to l_p . Moreover, if $1 < p < 2$ then L_p contains an uncomplemented subspace isomorphic to l_2 (certainly, this is obvious for $p = 1$). For $p > 2$ this result follows from [26] as it was noted in [27], p. 145. For $1 \leq p < 4/3$ this is a consequence of one W. Rudin's result [28]; on the whole interval $(1, 2)$ it was extended in [29]. At last, the existence in L_1 of an uncomplemented subspace isomorphic to l_1 was shown by J. Bourgain [30].

If $W \in \mathcal{A}^\sim((l_p)^F)$ then it is either $\mathcal{L}_p$ -space or $\mathcal{L}_2$ -space (in the sense of [31]) and, hence, contains a complemented subspace that is isomorphic to l_p (in the case $\mathcal{L}_p$; $1 \leq p < \infty$). It is easy to see that the only possibility is: $p > 2$ and $W \in (l_2)^F$. Indeed, in this case any space, which is isomorphic to the Hilbert space, has a complement in every space of the class $(l_p)^F$ that it contains, as it follows from [32].

In any other case there exists an isomorphic embedding u of W in the space $L_p(\mu)$ (for the suitable measure μ) such that the image uW has no complement in $L_p(\mu)$. $\square$

The result of M.I. Kadec and A. Pelczynski [32], mentioned above, was extended by B. Maurey [33] to the wide class of Banach spaces of type 2.,

In [33] it was shown:

- If a Banach space X is of type 2 and W is its subspace, isomorphic to the Hilbert space H , then there exists a projection $P : X \rightarrow W$ of norm

$$\|P\| \leq f(T_p(X)) d(W, H),$$

where the constant $f(T_p(X))$ depends only on $T_p(X)$.

An obvious consequence of this result is the following.

THEOREM 2. For any Banach space X of type 2 the class $\mathfrak{P}^\sim(X^F)$ is non-empty.

PROOF. As it follows from the aforementioned B. Maurey's theorem [33], the following inclusion is fulfilled: $(l_2)^F \subseteq \mathfrak{P}^\sim(X^F)$. $\square$

Notice that in the general (isometric) case classes $\mathcal{A}_0(X^f)$ and $\mathcal{E}(X^f)$ are different. E.g., if $X = l_p$ and $p \neq 2$ then the unique separable space in the class $\mathcal{E}((l_p)^f)$ is the space $L_p[0, 1]$. However, spaces l_p , $l_p \oplus_p L_p[0, 1]$ and a sequence of spaces $\{l_p^{(n)} \oplus_p L_p[0, 1] : n < \infty\}$ are pairwise non isometric and each of them belongs to $\mathcal{A}_0((l_p)^f)$. Notice, also, that in this case $\mathcal{A}_0((l_p)^f) = \mathcal{A}((l_p)^f)$ for all

$p \in [1, \infty]$ because of L_p contains an orthogonal complemented subspace l_2 only if $p = 2$.

THEOREM 3. *For every Banach space X classes $\mathcal{E}^\sim(X^F)$ and $\mathcal{A}^\sim(X^F)$ are identical.*

PROOF. The inclusion $\mathcal{E}^\sim(X^F) \subseteq \mathcal{A}^\sim(X^F)$ was noted before. To show the converse recall the following result, due to J. Stern [9]:

- *Let W, Z be a pair of Banach spaces; $W \hookrightarrow Z$. If W is an algebraic subspace of Z then W is a λ -reflecting subspace of Z for some $\lambda \leq 24$.*

Let $Z \in X^F$; $Y \in \mathcal{A}^\sim(X^F)$. Without loss of generality it may be assumed that $Y \in \mathfrak{q}\mathfrak{d}^\sim$ and $Y \hookrightarrow Z$. Hence, there exists such $\lambda < \infty$ that there exists a projection $P : Z \rightarrow Y$, $\|P\| \leq \lambda$.

Define on Z a new norm $\|\cdot\|'$ by taking as its unit ball the convex hull of the union of $\lambda^{-1}B(Y)$ and $B(Z)$ (recall that $B(X)$ denotes the unit ball of a Banach space X):

$$B(\langle Z, \|\cdot\|' \rangle) = \text{conv}\{(\lambda^{-1}B(Y)) \cup B(Z)\}.$$

Clearly, $\|y\|' = \|y\|$ for $y \in Y$; $\|z\|' \leq \lambda\|z\| \leq \lambda\|z\|'$ for $z \in Z$ and there exists a projection of norm one from $Z' = \langle Z, \|\cdot\|' \rangle$ onto Y .

Hence Z' may be represented as a direct orthogonal sum $Z' = Y \oplus W$, where W is C -finitely representable in Y for some $C < \infty$.

Let W be represented as a direct limit of the direct systems $(W_\alpha)_{\alpha \in A}$ of all its finite-dimensional subspaces: $W = \varinjlim W_\alpha$. Proceeding by induction on the directed set A , chose a set $(W'_\alpha)_{\alpha \in A}$ so that it also forms an isometric direct system, spaces $Y \oplus W'_\alpha$ are finitely representable in Y and $d(W_\alpha, W'_\alpha) \leq C'$ for some $C' < \infty$. Then the space $Z'' = \varinjlim Y \oplus W'_\alpha$ is finitely equivalent to Y , isomorphic to Z and Y is an algebraic subspace of Z'' . By the aforementioned Stern's theorem, Y is a λ -reflecting subspace of Z'' . Using the inverse isomorphisms we see that Y is λ' -reflecting subspace of Z . Since $Y \in \mathcal{A}^\sim(X^F)$ and $Z \in X^F$ are arbitrary, this yields that $\mathcal{A}^\sim(X^F) \subseteq \mathcal{E}^\sim(X^F)$. $\square$

Now it will be shown that the class $\mathfrak{P}^\sim(X^F)$ may be non-empty for a wide class of Banach spaces that are not of type 2. It will be needed the following result.

THEOREM 4. *Let a superreflexive class X^f contains a separable space E_X of almost universal disposition (with respect to $\mathfrak{M}(X^f)$). Let Z_1 and Z_2 be isomorphic subspaces of E_X of equal codimension (in E_X); $u : Z_1 \rightarrow Z_2$ be the corresponding isomorphism. Then there exists an isomorphic automorphism $\tilde{u} : E_X \rightarrow E_X$, which restriction to Z_1 is equal to u .*

PROOF. Let Z_1 and Z_2 be subspaces of E_X of equal codimension in E_X , i.e.

$$\text{codim}(Z_1) = \dim(E_X/Z_1) = \dim(E_X/Z_2) = \text{codim}(Z_2).$$

If $\text{codim}(Z_1) < \infty$ then assertions of the theorem are satisfied independently of a property of E_X to be a space of almost universal disposition. Indeed, for an arbitrary Banach space X any its subspaces of equal codimension, say, Y and Z , are isomorphic and each of them has a complement in X . So, X may be represented both as a direct sum $X = Y \oplus A$ and as $X = Z \oplus B$, where $\dim(A) = \dim(B) < \infty$.

Let $v : A \rightarrow B$ and $u : Y \rightarrow Z$ be isomorphisms. Clearly, $v \oplus u : X \rightarrow X$ is an automorphism of X , which extends u .

Assume that Z_1 and Z_2 are isomorphic subspaces of E_X of infinite codimension. Let $u : Z_1 \rightarrow Z_2$ be the corresponding isomorphism.

Let Z_1 be represented as a direct limit (= as the closure of union) of a chain

$$Z'_1 \hookrightarrow Z'_2 \hookrightarrow \dots \hookrightarrow Z'_n \hookrightarrow \dots \hookrightarrow Z_1;$$

namely, $Z_1 = \overline{\cup\{Z'_n : n < \omega\}}$ and $\cup Z'_n$ be dense in Z_1 . Clearly,

$$uZ'_1 \hookrightarrow uZ'_2 \hookrightarrow \dots \hookrightarrow uZ'_n \hookrightarrow \dots \hookrightarrow uZ_1 = Z_2$$

and the union $\cup uZ'_n$ is dense in Z_2 .

Let $(e_n)_{n < \omega}$ be a sequence of linearly independent elements of E_X of norm one, which linear span is dense in E_X .

Denote the restriction $u|_{Z'_n}$ by u_n and define by induction two sequences (f_n) and (g_n) of elements of E_X and a sequence (v_n) of isomorphisms.

Let f_1 be an element of $(e_n)_{n < \omega}$ with the minimal number, which does not belong to Z'_1 .

Put $W_1 = \text{span}(Z'_1 \cup \{f_1\})$ (recall that $\text{span}(A)$ denotes the *closure* of the linear span of A).

Let $\varepsilon > 0$ and $v_1 : W_1 \rightarrow E_X$ be an extension of u_1 such that

$$\|v_1\| \|v_1^{-1}\| \leq (1 + \varepsilon) \|u_1\| \|u_1^{-1}\|.$$

Put $g_1 = v_1 \circ f_1$; $U_1 = v_1 W_1$.

Let g_2 be an element of $(e_n)_{n < \omega}$ with the minimal number, which does not belong to $\text{span}(uZ'_1 \cup U_1)$.

Let $U_2 = \text{span}(uZ'_1 \cup U_1 \cup \{g_2\})$.

Since E_X is a space of almost universal disposition, there exists an isomorphism $(v_2)^{-1}$ which extends both $(u_2)^{-1}$ and $(v_1)^{-1}$ and satisfies the inequality

$$\|v_2\| \|v_2^{-1}\| \leq (1 + \varepsilon^2) \max\{\|u_2\| \|u_2^{-1}\|; \|v_1\| \|v_1^{-1}\|\}.$$

Let $f_2 = (v_2)^{-1} \circ g_2$. This close the first step of the induction.

Assume that $\{f_1, f_2, \dots, f_n\}$; $\{g_1, g_2, \dots, g_n\}$; $\{v_1, v_2, \dots, v_n\}$; $\{U_1, U_2, \dots, U_n\}$ and $\{W_1, W_2, \dots, W_n\}$ are already chosen. If n is odd, we choose sequentially:

f_{n+1} to be an element of $(e_n)_{n < \omega}$ with the minimal number, which does not belongs to Z'_{n+1} ;

$W_{n+1} = \text{span}(Z'_{n+1} \cup \{f_{n+1}\})$;

$v_{n+1} : W_{n+1} \rightarrow E_X$ be an extension of u_{n+1} such that

$$\|v_{n+1}\| \|v_{n+1}^{-1}\| \leq (1 + \varepsilon^{n+1}) \|u_{n+1}\| \|u_{n+1}^{-1}\|;$$

$g_{n+1} = v_{n+1} \circ f_{n+1}$; $U_{n+1} = v_{n+1} W_{n+1}$.

If n is even, we choose sequentially:

g_{n+1} to be an element of $(e_n)_{n < \omega}$ with the minimal number, which does not belong to $\text{span}(uZ'_n \cup U_n)$;

$U_{n+1} = \text{span}(uZ'_n \cup U_n \cup \{g_{n+1}\})$;

$(v_{n+1})^{-1}$, which extends both $(u_{n+1})^{-1}$ and $(v_n)^{-1}$ and satisfies

$$\|v_{n+1}\| \|v_{n+1}^{-1}\| \leq (1 + \varepsilon^2) \max\{\|u_{n+1}\| \|u_{n+1}^{-1}\|; \|v_n\| \|v_n^{-1}\|\};$$

$f_{n+1} = (v_{n+1})^{-1} \circ g_{n+1}$.

Since E_X is superreflexive, a sequence of isomorphisms (v_n) converges to an automorphism $V : E_X \rightarrow E_X$, which satisfies

$$\|V\| \|V^{-1}\| \leq \prod_{n=1}^{\infty} (1 + \varepsilon^n) \|u\| \|u^{-1}\|.$$

□

THEOREM 5. *Let X^f be a superreflexive class that contains a separable space E_X of almost universal disposition. Then $E_X \in \mathcal{E}^\sim(X^F) = \mathfrak{P}^\sim(X^F)$.*

PROOF. Let $Z \in X^F$; $j : E_X \rightarrow Z$ be an isomorphic embedding. Without loss of generality it may be assumed that Z is separable (if Z is non separable, it has a separable complemented subspace Z_0 , which contains jE_X). Also, it may be assumed that there exists an isometric embedding $k : Z \rightarrow E_X$ (since $E_X \in \mathcal{E}(X^f)$). Hence, the composition $k \circ j$ is an isomorphic embedding of E_X into E_X .

It may be assumed without the loss of generality that there exists an isometric embedding w of E_X into E_X .

There exists a bounded projection $P : E_X \rightarrow wE_X$. According to the previous theorem, there exists an automorphism $v : E_X \rightarrow E_X$, which sends jE_X to wE_X . Clearly, this implies that there exists a bounded projection from Z onto jE_X . Since Z is arbitrary, $E_X \in \mathfrak{P}^\sim(X^F)$. □

Analogously to $\mathfrak{P}^\sim(X^F)$ it may be defined the class $\mathfrak{J}^\sim(X^F)$.

DEFINITION 5. *Let X be a Banach space and $E <_F X$. E belongs to the class $\mathfrak{J}^\sim(X^F)$ provided for any pair $Y_0 \hookrightarrow Y$ where $Y <_F X$ every operator $u : Y_0 \rightarrow Y$ has a bounded extension $\tilde{u} : Y \rightarrow E$.*

THEOREM 6. *Let X^f be superreflexive and contains a separable space E_X of almost universal disposition. Let W be isomorphic to E_X . Then $W \in \mathfrak{J}^\sim(X^F)$.*

PROOF. Let W be isomorphic to E_X ; $u : E_X \rightarrow W$ be the corresponding isomorphism. Let $v : W \rightarrow E_X$ be the isomorphic embedding.

Let $Y_0 \hookrightarrow Y$; $Y <_F E_X$ and Y be separable. Let $i : Y_0 \rightarrow Y$ be an operator. Then

$$v \circ i : Y_0 \rightarrow vW \hookrightarrow E_X$$

and this mapping may be extended to the operator $V : Y \rightarrow E_X$ of the same norm.

Put $A = VY \cap vW$.

Since E_X is a space of almost universal disposition, there exists a subspace $W_0 \hookrightarrow E_X$ of infinite codimension such that $A \hookrightarrow W_0$ and $W_0 \in \mathcal{E}((E_X)^F)$. Hence, there exists an automorphism $U : E_X \rightarrow E_X$ such that $UW_0 = W$. Certainly, this isomorphism may be chosen to fix $(v \circ i)Y_0 \hookrightarrow E_X$.

Clearly, $U \circ V$ extends $v \circ i$ and, hence, $v^{-1} \circ U \circ V$ extends i . □

COROLLARY 1. *Let X^f be a superreflexive class that contains a separable space E_X of almost universal disposition. Let W be isomorphic to E_X . Then $W \in \mathfrak{P}^\sim(X^F)$.*

PROOF. As it was shown before, condition on X^f implies that $\mathfrak{P}(X^f) = \mathfrak{J}(X^f)$. Hence, by the previous theorem, $W \in \mathfrak{J}^\sim(X^F)$. Let $Z \in X_F$ and $W \hookrightarrow Z$. Then this identical embedding may be extended to the operator $P : Z \rightarrow Z$ such that $P|_W = Id_W$. Clearly, P is a projection. □

2. Bounded extension of operators. 2. A non-separable case.

Because of every class of crudely finite equivalence X^F , which contains a quotient-closed divisible class Y^f , contains also its Gurarii compression $\Gamma(X^f)$, and, as it was shown before, every superreflexive class of kind $\Gamma(X^f)$ contains a

separable space of almost universal disposition E_X with respect to $\mathfrak{M}(\Gamma(X^f))$, it follows that for every quotient-closed superreflexive divisible class X^f both classes $\mathfrak{P}^\sim(X^f)$ and $\mathfrak{J}^\sim(X^f)$ are equal and non-empty. However, it was proved the existence of only one member of these classes, namely - E_X . The existence of other spaces without assuming set theoretical hypothesis is doubtful.

Here it will be considered a non-separable case and it will be shown that under certain set-theoretical assumptions (e.g., under assumption of existence an inaccessible cardinal) classes $\mathfrak{P}^\sim(X^f)$ and $\mathfrak{J}^\sim(X^f)$ are confinal with X^f and are non-empty in the superreflexive case.

Recall some definitions on cardinal numbers.

Ordinals will be denoted by small Greece letters α, β, γ . *Cardinals* are identified with the least ordinals of given cardinality and are denoted by $\iota, \tau, \varkappa$ (may be, with indices). As usual, ω and ω_1 denote respectively the first infinite and the first uncountable cardinals (= ordinals).

A set of elements (of arbitrary nature) $\{a_\alpha : \alpha < \varkappa\}$ will be called *the $\varkappa$ -sequence*.

For a cardinal τ its *predecessor* (i.e., the least cardinal $\varkappa$, which is strongly greater than τ) is denoted by τ^+ .

The *confinality* of τ , $\text{cf}(\tau)$ is the least cardinality of a set $A \subset \tau$ such that $\tau = \sup A$.

Let A, B be sets. The symbol ${}^B A$ denotes the set of all functions from B to A .

In a general case, the cardinality of the set ${}^B A$ is denoted either by $\text{card}(A)^{\text{card}(B)}$ or by $\varkappa^\tau$, if $\text{card}(A) = \varkappa$; $\text{card}(B) = \tau$.

The symbol $\exp(\tau)$ (or, equivalently, 2^τ) denotes the cardinality of the set $\text{Pow}(\tau)$ of all subsets of τ .

A cardinal τ is said to be *regular* (resp., *singular*) if its confinality $\text{cf}(\tau) = \tau$ (resp., if $\text{cf}(\tau) < \tau$).

A cardinal ι is said to be *inaccessible* provided it is regular and $\exp(\varkappa) \leq \iota$ for every $\varkappa < \iota$.

The *continuum hypothesis* (CH) is the assumption

$$\omega^+ \stackrel{\text{def}}{=} \omega_1 = \exp(\omega).$$

The *general continuum hypothesis* (GCH) is the assumption

$$\tau^+ = \exp(\tau) \quad \text{for every infinite cardinal } \tau.$$

Let τ be a cardinal. Put

$$\varkappa^* = \sum \{\exp(\tau) : \tau < \varkappa\}$$

It is known that there are arbitrary large cardinals $\varkappa$ having the property

$$\varkappa = \varkappa^*.$$

In other words, a class of all cardinals with the property $\varkappa = \varkappa^*$ is confinal with the class of all cardinals. However, there known such models of the set theory, in which every cardinal $\varkappa$ with the property $\varkappa = \varkappa^*$ is singular.

From the other hand, in assuming the general continuum hypothesis (GCH) every regular cardinal $\varkappa$ has the property $\varkappa = \varkappa^*$. Certainly, any inaccessible cardinal ι is regular and enjoys the property $\iota = \iota^*$.

Recall that the existence of an inaccessible cardinal (as well as the CH or the GCH) cannot be proved in ZFC.

In what follows it will be assumed that there exists a regular cardinal κ with the property $\kappa = \kappa^*$. Every such cardinal will be called the **star-cardinal.**

Recall that a Banach space X is said to be an *envelope* (of the class X^f) if every Banach space Y , which is finitely representable in X and is of dimension $\dim Y \leq \dim X$ is isometric to a subspace of X . In [34] it was shown that for every cardinal κ with the property $\kappa = \kappa^*$ (it does not matter regular or singular) every class X^f of finite equivalence (generated by an infinite-dimensional space X) contains an envelope of this class of dimension κ .

It will be convenient to introduce one more definition.

DEFINITION 6. *A Banach space X is said to be f -saturated if X is a space of almost universal disposition with respect to a class $(X^{<f})_{\dim X}$ of all Banach spaces that are finitely representable in X and whose dimension is strictly less than $\dim X$;*

$$(X^{<f})_{\dim X} = \{Y <_f X : \dim Y < \dim X\}.$$

THEOREM 7. *Let X^f enjoys the isomorphic amalgamation property. Then for every **star-cardinal** κ the class X^f contains an f -saturated space $E_\kappa \in X^f$ of dimension κ .*

PROOF. Let F be an envelope of X^f of dimension κ . Let $(f_i)_{i < \kappa}$ be dense in the unit sphere of F .

For every $A \subset \kappa$ of cardinality $\text{card } A < \kappa$ put $F_A = \text{span}\{f_i : i \in A\}$. Surely, there are just κ different spaces of kind F_A .

For $A, B \subset \kappa$, $\max\{\text{card } A, \text{card } B\} < \kappa$ let $u_{A,B} : F_A \rightarrow F_B$ be an isomorphic embedding (if it exists). The set of all such embeddings is of cardinality $\tau \leq \kappa$.

Let $\varepsilon < 1$; $n \in \mathbb{N}$; $i_{A,n} : F_A \rightarrow F$ be an isomorphic embedding of norm $\|i_{A,n}\| \|i_{A,n}^{-1}\| \leq 1 + \varepsilon^n$. The set of all such embeddings is of cardinality that does not exceed κ as well.

So, the set T of all V -formations of kind $t = \langle F_A, F, F_B, i_{A,n}, u_{A,B} \rangle$ may be indexed by elements of κ . Let $T = \{t_\alpha : \alpha < \kappa\}$.

Construct by induction an κ -sequence of spaces $(E'_\alpha)_{\alpha < \kappa}$ as a chain

$$E'_0 \hookrightarrow E'_1 \hookrightarrow \dots E'_\alpha \hookrightarrow \dots$$

Put $E'_0 = F$. If $\alpha = \beta + 1$ then as E'_α it will be chosen an amalgam of the V -formation $\langle F_A, E'_\beta, F_B, v_{A,n}, u_{A,B} \rangle$, where $\langle F_A, F, F_B, i_{A,n}, u_{A,B} \rangle$ is just t_β in our indexation and $v_{A,n} = j \circ i_{A,n}$, where $j : F \rightarrow E'_\beta$ is a natural embedding.

If α is a limit ordinal (i.e. α has no representation $\alpha = \beta + 1$ for any β) put $E'_\alpha = \text{span}\{E'_\beta : \beta < \alpha\}$.

Put $E_{(1)} = \text{span}\{E'_\alpha : \alpha < \kappa\}$.

Now we repeat this inductive procedure κ times starting at the γ -th step with $E_{(\gamma)}$. Sequentially there will be obtained $E_{(1)} \hookrightarrow E_{(2)} \hookrightarrow \dots$

Put $E_\kappa = \text{span}\{E_{(\gamma)} : \gamma < \kappa\}$.

Certainly, $\dim E_\kappa = \kappa$. To show that E_κ is f -saturated chose a pair G, H of subspaces of E_κ , $\dim G \leq \dim H < \kappa$ and an isomorphic embedding $u : G \rightarrow H$. Since F is an envelope, G and H almost isometric to corresponding subspaces of F (say, F_G and F_H). Let $i : G \rightarrow E_\kappa$ be an isomorphic embedding, $\varepsilon > 0$ be fixed.

Since $\varkappa$ is regular, there exists such $\gamma < \varkappa$ that is $(1 + \varepsilon)$ -embedding of G into $E_{(\gamma)}$ and, hence, it may be extended to the embedding $\tilde{u}$ of B into $E_{(\gamma)}$ with $\|\tilde{u}\| \|\tilde{u}^{-1}\| \leq (1 + \varepsilon) \|u\| \|u^{-1}\|$.

Since G , H and $u : G \rightarrow H$ are arbitrary, this shown that $E_\varkappa$ is f -saturated. $\square$

According to [35] a Banach space X is said to be *subspace-homogeneous* if for every pair A, B of its isomorphic subspaces of equal (finite or infinite codimension) the isomorphism $v : A \rightarrow B$ may be extended to an isomorphical automorphism $\tilde{v}$ of X (i.e., $\tilde{v}|_A = v$). Using this terminology it may be said that the theorem 33 proves that every superreflexive separable space E of almost universal disposition (with respect to $\mathfrak{M}(E^f)$) is subspace-homogeneous. This theorem, that was proved firstly in the author's preprint [36], solves the problem of [35] on existence subspace-homogeneous Banach spaces that are differ from l_2 and c_0 .

The preceding theorem makes possible to prove the existence of non-separable subspace-homogeneous spaces (in assumption on existence of star-cardinals). The question on existence of non-separable subspace-homogeneous spaces also was posed in [35].

COROLLARY 2. *Let $E_\varkappa$ be a dual f -saturated space of the class X^f (this means that $E_\varkappa = W^*$ for some $W \in \mathcal{B}$). Then $E_\varkappa$ is subspace-homogeneous.*

In particular, $E_\varkappa \in \mathfrak{P}^\sim(X^F)$ ($= \mathfrak{J}^\sim(X^F)$)

PROOF. The proof literally repeats the proof of theorems 33 and 34 with obvious changes: in the present proof the transfinite induction is used.. At the last step of the proof is used the w^* -compactness of the unit ball of $E_\varkappa$. $\square$

THEOREM 8. *Let E be f -saturated and dual (i.e., $E = W^*$ for some $W \in \mathcal{B}$; in particular, E may be separable); F be a complemented subspace of E . Then $F \in \mathfrak{P}^\sim(X^F)$ ($= \mathfrak{J}^\sim(X^F)$).*

PROOF. Let A, B be isomorphic subspaces of E ; $A \hookrightarrow F \hookrightarrow E$; $B \hookrightarrow F \hookrightarrow E$, which are of equal (finite or infinite) codimension in F . Let $u : A \rightarrow B$ be the corresponding isomorphism. Let $U : E \rightarrow E$ be an automorphism of E that extends u ; $Ua = ua$ for all $a \in A$.

Let $P : E \rightarrow F$ be a projection. Certainly, $UF \hookrightarrow E$ is also a complemented subspace of E . Let $Q : E \rightarrow UF$ be the corresponding projection

So, $F \cap UF$ is also a complemented subspace of E , which contains B . Clearly, $F \cap UF$ is a complemented subspace of both F and UF . Let F' be its complement in F (resp., let F'' be its complement in UF). Clearly, F' and F'' are isomorphic. Let $v : F' \rightarrow F''$ be the corresponding isomorphism; id be identical on $F \cap UF$. Then the operator $v \circ id \circ U|_F$ is an automorphism of F , which restriction to A is equal to u . $\square$

REMARK 1. *In [36] it was shown that if E is subspace-homogeneous and its l_p -spectrum $S(E)$ contains a point $p \neq 2$ then E and $E \oplus E$ are non-isomorphic. So, from the preceding theorem it follows that there exists a pair of non-isomorphic separable spaces of $\mathfrak{P}^\sim(X^F)$ ($= \mathfrak{J}^\sim(X^F)$) when E satisfies the mentioned conditions.*

COROLLARY 3. *In assumption the GCH every superreflexive quotient-closed divisible class X^f contains spaces from $\mathfrak{P}^\sim(X^F)$ (and, thus, from $\mathfrak{J}^\sim(X^F)$) of arbitrary large cardinality.*

The same is true if we assume the existence of inaccessible cardinals.

Now it will be shown that the condition of regularity of $\varkappa$ cannot be omitted, at least, in the case when the class X^f is not super-stable in the sense of [37-38]. Recall the definition.

DEFINITION 7. ([37]). A Banach space X is said to be stable provided for any two sequences (x_n) and (y_m) of its elements and every pair of ultrafilter D, E over $\mathbb{N}$

$$\lim_{D(n)} \lim_{E(m)} \|x_n + y_m\| = \lim_{E(m)} \lim_{D(n)} \|x_n + y_m\|.$$

The notations $D(n)$ and $E(m)$ are used here (instead of D and E) to underline the variable (n or m respectively) in expressions like $\lim_{D(n)} f(n, m)$.

DEFINITION 8. ([38]). A Banach space X is said to be superstable if every its ultrapower $(X)_D$ is stable.

Let X be a Banach space.

A sequence $\{x_n : n < \infty\}$ of elements of X is said to be

- *Spreading*, if for any $n < \infty$, any $\varepsilon > 0$, any scalars $\{a_k : k < n\}$ and any choosing of $i_0 < i_1 < \dots < i_{n-1} < \dots$; $j_0 < j_1 < \dots < j_{n-1} < \dots$ of natural numbers

$$\left\| \sum_{k < n} a_k x_{i_k} \right\| = \left\| \sum_{k < n} a_k x_{j_k} \right\|.$$

- *Symmetric*, if for any $n < \omega$, any finite subset $I \subset \mathbb{N}$ of cardinality n , any rearrangement ς of elements of I and any scalars $\{a_i : i \in I\}$,

$$\left\| \sum_{i \in I} a_i z_i \right\| = \left\| \sum_{i \in I} a_{\varsigma(i)} z_i \right\|.$$

DEFINITION 9. Let X be a Banach space. Its IS-spectrum $IS(X)$ is a set of all (separable) spaces $\langle Y, (y_i) \rangle$ with a spreading basis (y_i) which are finitely representable in X .

THEOREM 9. A class X^f is superstable if and only if every member $\langle Y, (y_i) \rangle$ of its IS-spectrum has a symmetric basis.

PROOF. The first part of the theorem was proved in [37].

Conversely, let every $\langle Z, (z_i) \rangle$ has a symmetric basis. Suppose that X is not superstable. Then there exists a space from X^f which is not stable (it may be assumed that X is not stable itself). By [37] there are such sequences (x_n) and (y_m) of elements of X that

$$\sup_{m < n} \|x_n + y_m\| > \inf_{m > n} \|x_n + y_m\|.$$

Let D be a countably incomplete ultrafilter over $\mathbb{N}$. Put

$$X_0 \stackrel{\text{def}}{=} X; \quad X_n \stackrel{\text{def}}{=} (X_{n-1})_D; \quad n = 1, 2, \dots; \quad X_\infty = \overline{\bigcup_{n \geq 1} X_n}.$$

Here is assumed that X_n is a subspace of $X_{n+1} = (X_n)_D$ under the canonical embedding $d_{X_n} : X_n \rightarrow (X_n)_D$.

Let D, E be ultrafilters over $\mathbb{N}$. Their product $D \times E$ is a set of all subsets A of $\mathbb{N} \times \mathbb{N}$ that are given by

$$\{j \in \mathbb{N} : \{i \in \mathbb{N} : (i, j) \in A\} \in D\} \in E.$$

Certainly, $D \times E$ is an ultrafilter and for every Banach space Z the ultrapower $(Z)_{D \times E}$ may be in a natural way identified with $((Z)_D)_E$.

So, the sequence $(x_n) \subset X$ defines elements

$$\begin{aligned} \mathfrak{x}_1 &= (x_n)_D \in (X)_D; \\ \mathfrak{x}_2 &= (x_n)_{D \times D} \in ((X)_D)_D; \\ &\dots \dots \dots \dots \dots \dots \\ \mathfrak{x}_k &= (x_n) \underbrace{D \times D \times \dots \times D}_{k \text{ times}} \in \underbrace{(((X)_D)_D \dots)_D}_{k \text{ times}}; \\ &\dots \dots \dots \dots \dots \dots \end{aligned}$$

Notice that $\mathfrak{x}_k \in X_k \setminus X_{k-1}$. It is easy to verify that $(\mathfrak{x}_k)_{k < \infty} \subset X_\infty$ is a spreading sequence. Since $X_\infty \in X^f$, it is symmetric. Moreover, for any $z \in X$, where X is regarded as a subspace of X_∞ under the direct limit of compositions

$$d_{X_n} \circ d_{X_{n-1}} \circ \dots \circ d_{X_0} : X \rightarrow X_n,$$

the following equality is satisfied: for any pair $m, n \in \mathbb{N}$

$$\|x_n + z\| = \|x_m + z\|.$$

Since (x_n) and z are arbitrary elements of X , this contradicts with the inequality $\sup_{m < n} \|x_n + y_m\| > \inf_{m > n} \|x_n + y_m\|$. $\square$

THEOREM 10. *Let E be an envelope of the class E^f , which is not super-stable. Let $\dim E = \varkappa$ and $\varkappa$ is singular. Then E is not f -saturated.*

PROOF. Since E^f is not super-stable, there exists a space $\langle W, (w_n) \rangle$ with a spreading basis, which is not symmetric one. Consider a vector space $c_{00}(\varkappa)$ of all $\varkappa$ -sequences of reals all but finitely many members of which are vanished.

Let $e_\alpha = (\delta_{\alpha\beta})_{\beta < \varkappa} \in c_{00}(\varkappa)$ ($\alpha < \varkappa$), where $\delta_{\alpha\beta}$ be the generalized Kronecker symbol; $\delta_{\alpha\beta} = 0$ when $\alpha \neq \beta$; $\delta_{\alpha\beta} = 1$ when $\alpha = \beta$.

Every element $x \in c_{00}(\varkappa)$ is of kind $x_A = \sum \{x_\alpha e_\alpha : \alpha \in A\}$, where $A \subset \varkappa$ is finite.

Let $A = \{\alpha_0, \alpha_1, \dots, \alpha_m\}$ and $\alpha_0 < \alpha_1 < \dots < \alpha_m$. Define on $c_{00}(\varkappa)$ a norm, which is given by

$$\|x_A\| = \left\| \sum_{i=0}^m x_{\alpha_i} w_i \right\|_W.$$

Let $W(\varkappa)$ be a completion of $c_{00}(\varkappa)$ under this norm.

Surely, $W(\varkappa)$ has a spreading, non symmetric basis, is finitely representable in $E_\varkappa$ and is isometric to a subspace of $E_\varkappa$ (since $\dim W(\varkappa) = \varkappa = \dim E_\varkappa$ and $E_\varkappa$ is an envelope).

Since $\varkappa$ is singular, $\text{cf}(\varkappa) = \tau < \varkappa$. Let $(\varkappa_\gamma)_{\gamma < \tau}$ be an increasing τ -sequence of cardinals (that are less than $\varkappa$), such that $\sup\{\varkappa_\gamma : \gamma < \tau\}$

Consider two subspaces of $W(\varkappa)$, say $W_1(\tau)$ and $W_2(\tau)$.

$W_1(\tau)$ is spanned by the first τ elements of e_α ;

$$W_1(\tau) = \text{span}\{e_\alpha : \alpha < \tau\}.$$

$W_2(\tau)$ is spanned by those e_α 's, whose indices belong to the set $(\varkappa_\gamma)_{\gamma < \tau}$;

$$W_2(\tau) = \text{span}\{e_{\tau_\beta} : \beta < \tau\}.$$

Surely, $W_1(\tau)$ and $W_2(\tau)$ are isometric, but no automorphism $u : E \rightarrow E$ sends $W_1(\tau)$ to $W_2(\tau)$. $\square$

3. References

- (1) Nachbin I. *A theorem of Hahn-Banach type for linear transformations*, Trans. AMS **68** (1950) 28-46
- (2) Goodner D. A. *Projections in normed and linear spaces*, Trans. AMS **69** (1950) 89-108
- (3) Kelley J. *Banach spaces with the extension property*, Trans. AMS **72** (1952) 323-326
- (4) Lindenstrauss J. *On a problem of Nachbin concerning extension of operators*, Israel J. Math. **2:1** (1963) 75-90
- (5) Lindenstrauss J. *Extension of compact operators*, Memoirs AMS **48** (1964)
- (6) Tokarev E.V. *Injective Banach spaces in the finite equivalence classes* (transl. from Russian), Ukrainian Mathematical Journal **39:6** (1987) 614-619
- (7) Dacunha-Castelle D., Krivine J. -L. *Applications des ultraproducts à l'étude des espaces et des algèbres de Banach*, Studia Math. **41** (1972) 315 - 334
- (8) Schwartz L. *Geometry and probability in Banach spaces*, Bull. AMS **4:2** (1981) 135-141
- (9) Stern J. *Ultrapowers and local properties in Banach spaces*, Trans. AMS **240** (1978) 231-252
- (10) Yasuhara M. *The amalgamation property, the universal homogeneous models and generic models*, Math. Scand. **34** (1974) 5 - 36
- (11) Lindenstrauss J., Rosenthal H.P. *The $\mathcal{L}_p$ -spaces*, Israel J. Math. **7** (1969) 325-349
- (12) Davis W.J., Figiel T., Johnson W.B., Pełczyński A. *Factoring weakly compact operators*, J. Funct. Anal. **17** (1974) 311-327
- (13) Heinrich S. *Ultraproducts in Banach space theory*, J. Reine Angew. Math. **313** (1980) 72-104
- (14) Gurarii V.I. *On inclinations and spreadings of subspaces of Banach spaces* (in Russian), Teor. Funct., Funct. Analysis and Appl. **1** (1965) 194-204
- (15) Pisier G. *On the duality between type and cotype*, Lect. Notes in Math. **939** (1982)
- (16) Pietsch A. *Absolut p -summierende Abbildungen in normierten Räumen*, Studia Math. **28** (1967) 333-353
- (17) Pełczyński A. *p -integral operators commuting with group representations and examples of quasi p -integral operators, which are not p -integral*, Studia Math. **33** (1969) 63-70
- (18) Pisier G. *Counterexample to a conjecture of Grothendieck*, Acta Math. **151** (1983) 181-208
- (19) Grothendieck A. *Sur quelques points d'algèbre homologique*, Tôhoku Math. J., second series, **9:2,3** (1957) 119-221
- (20) Heinrich S. *Finite representability and super - ideals of operators*, Diss. Math. (Rozprawy Matematyczne) **172** (1980) Warszawa
- (21) Gurarii V.I. *Spaces of universal disposition, isotropic spaces and the Mazur problem on rotations in Banach spaces*, Sibirsk. Mat. Journ. (in Russian) **7** (1966) 1002-1013
- (22) Maurey B., Pisier G. *Séries de variables aléatoires vectorielles indépendantes et propriétés géométriques des espaces de Banach*, Studia Math. **58** (1976) 45-90

- (23) Rosenthal H.P. *On a theorem of J. -L. Krivine concerning block finite representation of l_p in general Banach spaces*, J. Funct. Anal. **25** (1978) 197-225
- (24) Stromquist W. *The maximum distance between two-dimensional spaces*, Math. Scand. **48** (1981) 205-225
- (25) Kadec M.I., Snobar L.G. *On some functionals on Minkowski's spaces* (in Russian), Mat. Zametki **10** (1971) 453-458
- (26) Rosenthal H.P. *Projections onto translation-invariant subspaces of $L_p(G)$* , Memoirs AMS **63** (1968) 1-84
- (27) Lindenstrauss J. and Tzafriri L. *Classical Banach spaces*, Lecture Notes in Math. **338** (1973) 1-242
- (28) Rudin W. *Trigonometric series with gaps*, J. Math. Mech. **9** (1960) 203-227
- (29) Bennet G., Dor L.E., Gudman V., Johnson W.B., Newmann C.M. *On uncomplemented subspaces of L_p , $1 < p < 2$* , Israel J. Math. **26** (1977) 178-187
- (30) Bourgain J. *Complementation de sous-espaces L_1 dans les espaces L_1* , Sem. Funct. Anal. École Polytech. Palaiseau, exp. **27** (1980) 1-7
- (31) Lindenstrauss J., Pełczyński A. *Absolutely summing operators in $\mathcal{L}_p$ -spaces and their applications*, Studia Math. **29** (1968) 275-326
- (32) Kadec M.I. and Pełczyński A. *Bases, lacunary sequences and complemented subspaces in the spaces L_p* Studia Math. **21** (1962) 161-176
- (33) Maurey B. *Quelques problèmes de factorization d'opérateurs linéaires*, Actes du Congress Int. Math. Vancouver, 1974 v. **2** (1975) 75-79
- (34) Tokarev E.V. *Problem of envelopes of locally convex spaces* (transl. from Russian), Siberian Mathematical Journal **31:1** (1990) 173-175
- (35) Lindenstrauss J., Rosenthal H.P. *Automorphisms in c_0 , ℓ_1 and m* , Israel J. Math. **7** (1969) 227-239
- (36) Tokarev E.V. *A solution of one problem of Lindenstrauss and Rosenthal on subspace-homogeneous and quotient-homogeneous Banach spaces with application to the approximation problem and to the Schroeder - Bernstein problem*, Preprint. Printed electronically in **www.ArXiv.org** at June, 03, 2002, **math.FA/0206013**
- (37) Krivine J. -L., Maurey B. *Espaces de Banach stables*, Israel J. Math. **39** (1981) 273-295
- (38) Raynaud Y. *Espaces de Banach superstables*, C. R. Acad. Sci. Paris, Sér. A. **292:14** (1981) 671-673